


Should *cure* be the goal for multiple myeloma?



Brought to you by SES




Scientific Educαtion Support




Summary and conclusions

Sagar Lonial

Winship Cancer Institute of Emory University, Atlanta, US



Experience from the RVD 1000 series: a new benchmark for PFS and OS




CI, confidence interval; No, number; OS, overall survival; PFS, progression-free survival; RVD, lenalidomide, bortezomib, dexamethasone. Joseph NS, et al. *J Clin Oncol*. 2020;38(17):1928-1937

'Cure' is in the eye of the beholder

- A complicated disease like MM is going to have many different definitions of cure.
- These may not fully identify with existing definitions, as this is not HL, testicular CA, DLBCL, or ALL.
- While it is the hope that early intervention with precursor conditions may increase the cure rate, this is not proven, and evidence suggests this may not be the case.
- The use of MRD as a surrogate for cure may not be ready for prime time;
 - Sustained MRD negativity may be more meaningful.
- Need to distinguish operational or functional cure from old definitions or standards for cure.
- Do not de-escalate therapy till we are clear that outcomes are BETTER (the same is not a step forward).
- While surrogates are great, they must be demonstrated to have better PFS and OS to be meaningful.

Model for elimination of the malignant clone



Consolidation/maintenance (to achieve cure)



